Athedround A

COLLEGE PHYSICS

ELECTRICITY AND MAGNETISM, LIGHT, AND ATOMIC PHYSICS

ADDISON-WESLEY PUBLISHING COMPANY, INC.

FRANCIS WESTON SBARS

Dartmouth College

gud,

MARK W. ZEMANSKY

The City College of New York

TRIBD EDITION

ADDISON-WESLEY PUBLISHING COMPANY, INC. READING, MABGACHUSETTS, U.S.A. LONDON, ENGLAND

all rights reserved. This book, or parts thereof, may not de Reproduced in any form with-Library of Congress Catalog Card No. 60-5014 OUT WRITTEN PERHISSION OF THE PUBLISHER, Printed in the United States of America Third Edition

BEST AVAILABLE COPY



BEST AVAILABLE COPY

Potal internal reflection can occur only when a ray is incident on the surface of a medium whose index is smaller than that of the medium in upper medium but is totally internally reflected at the boundary surface. TOTAL INTERNAL REFLECTION which the ray is traveling.

The critical angle for two given substances may be found by setting $\phi' = 90^{\circ}$ or $\sin \phi' = 1$ in Snell's law. We then have

$$\sin \phi_c = \frac{n'}{n}. \tag{40-3}$$

The critical angle of an air-glass surface, taking 1.50 as a typical index of refraction of glass, is

$$\sin \phi_c = \frac{1}{1.50} = 0.67, \quad \phi_c = 4$$

Pic. 40-5. (a) A train of plane waves is in part reflected and in part refracted at the boundary between two media. (b) The waves in (a) are represented by rays. (c) For simplicity, only a single incident, reflected, and

Referenced waves


Refracted 133

દ

3

45-90° as totally reflecting surfaces. The advantages of totally reflecting This angle, very conveniently, is slightly less than 45°, which makes possible the use in many optical instruments of prisms of angles 45°prisms over metallic surfaces as reflectors are, first, that the light is totally reflected, while no metallic surface reflects 100% of the light incident on it, and second, the reflecting properties are permanent and not affected by tarnishing. Offsetting these is the fact that there is some loss of light by reflection at the surfaces where light enters and leaves the prism, atthough recently discovered methods of coating the surfaces with so-called "nonreflecting" films can reduce this loss considerably.

The simplest type of reflecting prism is shown in Fig. 40-7. Its angles are 46"-45"-90°. Light incident normally on one of the shorter faces



diverging from a point source P in a medium of index n and striking the

urface of a second medium of index n', where n > n'. From Snell's law,

 $\sin \phi' = \frac{\pi}{\pi'} \sin \phi$.

40-4 Total internal reflection. Figure 40-6 shows a number of rays

lently equals unity (i.e., $\phi' = 90^{\circ}$) for some angle ϕ less than 90°. This is illustrated by ray 3 in the diagram, which emerges just grazing the

Since n/m' is greater than unity, sin o' is larger than sin o and evi-

angle and is designated by \$\psi_c\$ in the diagram. If the angle of incidence is greater than the critical angle, the sine of the angle of refraction, as

computed by Snell's law, is greater than unity. This may be interpreted to mean that beyond the critical angle the ray does not pass into the

the refracted ray emerges tangent to the surface is called the critical

surface at an angle of refraction of 90°. The angle of incidence for which

Fio. 40-6. Total internal reflection. critical angle.

Fra. 40-7. A totally reflecting

The angle of incidence ϕ_{ei} for which the angle of refraction is 90°, is called the

REPLECTION AND REPRACTION AT PLANE SURFACES [CHAP. 40]

Reflected

Incident

Reflected

ment A page PAGE 13/13 * RCVD AT 6/30/2005 4:13:55 PM [Eastern Daylight Time] * SVR:USPTO-EFXRF-1/0 * DNIS:8729306 * CSID:9149453281 * DURATION (mm-ss):03-163E . 13 ***

of incidence. The incident ray, the reflected ray, and the normal to the surface

When a ray of light is reflected, the angle of reflection is equal to the angle

To summarize the laws of reflection and refraction in terms of rays:

the reverse is true and the ray is bent away from the normal

bent toward the normal. If the light is traveling in the opposite direction,

refracted ray are drawn.

he refracted ray, and the normal to the surface at the point of incidence, all

is in the same plane.

When a ray of light is refracted, n sin $\phi = n' \sin \phi'$. The incident ray,

at the point of incidence, all lie in the same plane.